



1/20

DeathTRAK Homogeneous Cytotoxicity Assay












30 Lysis of PC-3 Measured by DeathTRAK Homogenous Assay Effect of Anti-Factor I Antibody on Complement-Mediated 10 20 [Monoclonal Antibody] (nanomolar) Fig. 5 FLU/Second rO

30 Effect of Anti-Factor I Antibody on Complement-Mediated Lysis of PC-3: Single-Point Readout Signal after 2.6 minutes [Monoclonal Antibody] (nanomolar) Fig. 6 009 S Brn 0 07/9


Effect of Additional ADP on Homogeneous DeathTRAK Signal









DeathTRAK Signal from Alive vs. Dead Cells (E. coli)





Protection of G3PDH in the Presence of ---3 mM DTT + 1% PICguws $-\Delta$ 3 mM DTT + 2% PICguws Dead PC-3 Cells Time (Hours) → Nothing Fig. 11 1200 _¬ BLU\Second 800 11/20






Effects of Three Antibiotics on Proliferation/Viability of E. coli K1 Measured by DeathTRAK






50 (Gram Positives with Three Antibiotics) DeathTRAK Toxicity Readout [Antibiotic] (µg/mL) Fig. 16 → Carbenicillin -∆-Vancomycin --- Gentamicin RLU/Second 50 10 16/20

Viability of Group-A Streptococcus Challenged with Three Antibiotics Measured by DeathTRAK



18/20

PhosTRAK Homogeneous Phosphatase Assay



Detection of Free Phosphate by PhosTRAK 400 pmol (Pi) Fig. 19 200 RLU\Second 100 19/20

